


REVOIS® PRO PATIENT CASES

Dr. Frederic Hermann M.Sc.




Case 1

Initial situation



Implantplaning



Implant insertion with surgical guides




Radiographic control a surgery






1 year follow-up




5 year follow-up






Case 2

Initial situation, regio 47




3D-Implantplaning



Situation after integration of the crown occlusally screw-fixed



Radiographie control 1 year after surgery




Source: Dr. Frederic Hermann M.Sc. - TEAM 15 - Praxis für Zahnmedizin - Poststrasse 15 - CH-6300 Zug - Switzerland

Radiographie control 3 year after surgery



Radiographie control 5 year after surgery



Case 3

Initial situation, regio 11-13



Initial situation, regio 11-13



Indication to extract tooth 11 and 13



Minimally invasive tooth extraction & immediate implantation




Minimally invasive tooth extraction & immediate implantation




Source: Dr. Frederic Hermann M.Sc. - TEAM 15 - Praxis für Zahnmedizin - Poststrasse 15 - CH-6300 Zug - Switzerland


Minimally invasive tooth extraction & immediate implantation



Minimally invasive tooth extraction & immediate implantation




Radiographie control after surgery




Situation at the end of the healing phase 3 month later



Situation at the end of the healing phase 3 month later



Open impression-faking



Open impression-faking



Abutments` fry-on



Scaffold try-on



Integration of bridge 3 month after implant insertion



Integration of bridge 3 month after implant insertion



Situation at 5 years follow-up



Radiographie control (for maxillary therapy planing) at 5 years follow-up

